

Bulletin of the British Myriapod Group 8 (1992)

A COLLECTION OF IRISH MYRIAPODS

R.E.Jones
14 Post Office Road, Dersingham, Norfolk, PE31 6HP

In March 1991 I made a visit to Southern Ireland during which I was able to collect various invertebrates. The results of the trip with regard to myriapods are set out below. The main collection was from Kerry but there were others which were made to and from the boat. One millipede and one centipede were new for Ireland (marked *). The name of the place (or the nearest to it on a map), the vice-county, the map reference and date form the heading for each site.

| | | |
|---|---------------------------------|---------|
| Killane, Wexford | T1309 | 22.3.91 |
| <i>Blaniulus guttulatus</i> | <i>Geophilus electricus</i> | |
| <i>Brachydesmus superus</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Polydesmus gallicus</i> | | |
| Granny Castle, Waterford | S5714 | 22.3.91 |
| | <i>Lithobius forficatus</i> | |
| | <i>L.microps</i> | |
| West of Piltown, Kilkenny | S4322 | 22.3.91 |
| <i>Glomeris marginata</i> | | |
| Clonmel, S.Tipperary | S1822 | 22.3.91 |
| | <i>Lithobius microps</i> | |
| Parking place just W.of Boolakennedy, R9319 | | 22.3.91 |
| S.Tipperary | | |
| <i>Brachydesmus superus</i> | <i>Lithobius forficatus</i> | |
| <i>Brachyiulus pusillus</i> | <i>L.microps</i> | |
| <i>Polydesmus gallicus</i> | | |
| Rathbarry, W.Cork | W3334 | 22.3.91 |
| <i>Polydesmus gallicus</i> | <i>Lithobius forficatus</i> | |
| | <i>L.variegatus</i> | |
| Rosscarbery Bay, W.Cork | W3234 | 23.3.91 |
| <i>Cylindroiulus latestriatus</i> | <i>Haplophilus subterraneus</i> | |
| <i>Glomeris marginata</i> | <i>Cryptops hortensis</i> | |
| <i>Ophyiulus pilosus</i> | <i>Lithobius melanops</i> | |
| <i>Polydesmus angustus</i> | <i>L.variegatus</i> | |

| | | | |
|-----------------------------------|--|----------------------------------|---------|
| Lough Hyne, W.Cork | | W0929 | 23.3.91 |
| <i>Brachydesmus superus</i> | | <i>Brachygeophilus truncorum</i> | |
| <i>Glomeris marginata</i> | | <i>Geophilus electricus</i> | |
| <i>Ophyiulus pilosus</i> | | <i>Necrophloeophagus flavus</i> | |
| <i>Polydesmus angustus</i> | | <i>Lithobius forficatus</i> | |
| <i>Proteroiulus fuscus</i> | | <i>L.borealis</i> | |
| Baltimore, W.Cork | | W04326 | 23.3.91 |
| <i>Brachychaeteuma melanops</i> | | <i>Cryptops hortensis</i> | |
| <i>Cylindroiulus punctatus</i> | | <i>Necrophloeophagus flavus</i> | |
| <i>Ophiodesmus albonanus</i> | | <i>Lithobius forficatus</i> | |
| Glengarriff, W.Cork | | V9157 | 24.3.91 |
| <i>Cylindroiulus punctatus</i> | | <i>Brachygeophilus truncorum</i> | |
| | | <i>Geophilus electricus</i> | |
| | | <i>Necrophloeophagus flavus</i> | |
| | | <i>Lithobius variegatus</i> | |
| N71 just over the border, S.Kerry | | V9060 | 24.3.91 |
| | | <i>Brachgeophilus truncorum</i> | |
| Moll's Gap, S.Kerry | | V8677 | 24.3.91 |
| | | <i>Brachygeophilus truncorum</i> | |
| | | <i>Geophilus insculptus</i> | |
| Island in the Upper Lake, S.Kerry | | | 23.3.91 |
| <i>Cylindroiulus punctatus</i> | | <i>Geophilus electricus</i> | |
| <i>Proteroiulus fuscus</i> | | <i>Necrophloeophagus flavus</i> | |
| | | <i>Schendyla nemorensis</i> | |
| | | <i>Lithobius borealis</i> | |
| Blackstones Bridge, S.Kerry | | V7186 | 25.3.91 |
| <i>Cylindroiulus punctatus</i> | | <i>Brachygeophilus truncorum</i> | |
| <i>Polydesmus angustus</i> | | <i>Cryptops hortensis</i> | |
| <i>Proteroiuls fuscus</i> | | <i>Lithobius borealis</i> | |
| | | <i>L.variegatus</i> | |
| Near Nedranone, S.Kerry | | V6059 | 26.3.91 |
| <i>Cylindroiulus latestriatus</i> | | <i>Brachygeophilus truncorum</i> | |
| | | <i>Geophilus carpophagus</i> | |
| | | <i>Schendyla nemorensis</i> | |
| | | <i>Lithobius borealis</i> | |
| Derrynane National Park, S.Kerry | | V5158 | 26.3.91 |
| <i>Cylindroiulus latestriatus</i> | | <i>Brachygeophilus truncorum</i> | |
| <i>Glomeris marginata</i> | | <i>Haplophilus subterraneus</i> | |
| <i>Ophyiulus pilosus</i> | | <i>Necrophloeophagus flavus</i> | |
| <i>Macrosternodesmus palicola</i> | | <i>Lithobius forficatus</i> | |
| <i>Polydesmus angustus</i> | | | |

| | | |
|-----------------------------------|----------------------------------|---------|
| Derrynane National Park, S.Kerry | V5258 | 26.3.91 |
| <i>Cylindroiulus punctatus</i> | <i>Geophilus electricus</i> | |
| <i>Glomeris marginata</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Opyiulus pilosus</i> | | |
| <i>Proteroius fuscus</i> | | |
| Near Cappamore, S.Kerry | V5664 | 26.3.91 |
| * <i>Chordeuma proximum</i> | <i>Brachygeophilus truncorum</i> | |
| | <i>Necrophloeophagus flavus</i> | |
| | <i>Lithobius variegatus</i> | |
| Valencia Island, S.Kerry | V3977 | 27.3.91 |
| <i>Brachydesmus superus</i> | <i>Brachygeophilus truncorum</i> | |
| | <i>Geophilus electricus</i> | |
| | <i>Lithobius forficatus</i> | |
| Valencia Island, S.Kerry | V4176 | 27.3.91 |
| <i>Cylindroiulus punctatus</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Ophyiulus pilosus</i> | | |
| <i>Proteroius fuscus</i> | | |
| Ballinskelligs, S.Kerry | V4366 | 27.3.91 |
| <i>Cylindroiulus latestriatus</i> | | |
| West of Templenoe, S.Kerry | V8269 | 28.3.91 |
| <i>Ophyiulus pilosus</i> | <i>Brachygeophilus truncorum</i> | |
| | <i>Lithobius variegatus</i> | |
| Parknasilla, S.Kerry | V7265 | 28.3.91 |
| | <i>Brachygeophilus truncorum</i> | |
| | <i>Geophilus electricus</i> | |
| | <i>Schendyla nemorensis</i> | |
| | <i>Lithobius melanops</i> | |
| | <i>L.variegatus</i> | |
| Gougane Barra Forest Park, W.Cork | W0765 | 28.3.91 |
| <i>Cylindroiulus punctatus</i> | <i>Lithobius borealis</i> | |
| | <i>L.variegatus</i> | |
| West of Kilgarvan, N.Kerry | V9972 | 28.3.91 |
| <i>Blaniulus guttulatus</i> | <i>Brachygeophilus truncorum</i> | |
| <i>Brachydesmus superus</i> | * <i>Henia brevis</i> | |
| <i>Macrosternodesmus palicola</i> | <i>Geophilus electricus</i> | |
| <i>Ophiodesmus albonanus</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Ophyiulus pilosus</i> | <i>Lithobius melanops</i> | |
| <i>Polydesmus angustus</i> | | |
| Rathmore, N.Kerry | W1493 | 29.3.91 |
| <i>Cylindroiulus punctatus</i> | <i>Geophilus insculptus</i> | |
| Near Lombardstown, Mid Cork | W4597 | 29.3.91 |
| | <i>Lithobius variegatus</i> | |

| | | |
|--|---------------------------------|---------|
| Near Clondulane on the Blackwater R. W8799 E.Cork | | 29.3.91 |
| <i>Brachydesmus superus</i> | <i>Schendyla nemorensis</i> | |
| <i>Brachyiulus pusillus</i> | | |
| <i>Cylindroiulus punctatus</i> | | |
| <i>Ophyiulus pilosus</i> | | |
| East of Ballyduff, Waterford | W9999 | 29.3.91 |
| <i>Ophyiulus pilosus</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Polydesmus gallicus</i> | <i>Lithobius forficatus</i> | |
| | <i>L.variegatus</i> | |
| River Barrow near New Ross, Kilkenny | S6824 | 29.3.91 |
| | <i>Cryptops hortensis</i> | |
| | <i>Lithobius variegatus</i> | |
| Arthurstown quay, Wexford | S7110 | 29.3.91 |
| <i>Cylindroiulus latestriatus</i> | <i>Necrophloeophagus flavus</i> | |
| <i>Polydesmus angustus</i> | <i>Strigamia maritima</i> | |
| <i>Tachypodoiulus niger</i> | | |

DISCUSSION

The two *Chordeuma proximum* males were collected from a ditch at the foot of a steep slope. The slope was wooded but there was nothing to suggest that it was in any way special; I think that *C.proximum* will be found to be not uncommon when it is looked for. *Henia brevis* came from the garden of a 19th century farmhouse. It was not recognised until it was back in England so the exact site is uncertain. It may be quite widespread or it may have been introduced with plants into the garden.

The *Brachychaeteuma* presents difficulties as it is a female but it has five ocelli on one side so the assumption is that it is *melanops*. The only other Irish specimen was also a female, collected by me in 1982 from Castletownshend in an adjacent grid square.

Macrosternodesmus palicola and *Ophiodesmus albonanus* were both found twice despite not being searched for in any special way. Looking under large stones produced the *O.albonanus*. Again, I think they will both prove to be common and widespread in Ireland. *Geophilus electricus* was found on a number of occasions, which shows it to be common. *Necrophloeophagus flavus* lived up to its reputation of being the commonest geophilomorph whilst *G.carpophagus* appeared to be uncommon.

Ireland is still little known from the myriapod point of view and the full list of species is undoubtably incomplete.